



ELSEVIER

Atmospheric Research 37 (1995) 119–131

ATMOSPHERIC
RESEARCH

Some observations on climate variability as seen in daily temperature structure

D.A. Robinson ^a, D.J. Leathers ^b, M.A. Palecki ^c, K.F. Dewey ^c

^a *Department of Geography, Rutgers University, New Brunswick, NJ 08903, USA*

^b *Center for Climatic Research, Department of Geography, University of Delaware, Newark, DE 19716, USA*

^c *Department of Geography, University of Nebraska, Lincoln, NE 68588, USA*

Accepted 17 November 1994

Abstract

Distinctive annual cycles of daily maximum and minimum temperature and temperature range are observed across the United States when averaging daily data for multiple decades. In some cases, abrupt changes in the range between daily maximum and minimum temperatures are observed within annual cycles. Such discontinuities, or steps, of 5°C may occur within a week, despite the fact that the daily data are averaged over several decades. Discontinuities are most pronounced in the fall and spring, and display differing characteristics of timing, rapidity, magnitude and direction of change across the nation. In northern and central portions of the eastern U.S., a winter minimum and summer maximum in temperature range is observed, while in the southern U.S. the opposite is found. Abrupt discontinuities are observed in early November in the Northeast and in late spring and early summer in the Southeast and Southwest.

Differences in mean daily temperature cycles between various portions of this century are a function of variations in maximum and minimum temperatures. The central U.S. had higher maximum and minimum temperatures during the 1930's than at other times this century, with little effect on daily temperature range. In the Northeast, the temperature range was lower in the 1930's, as a result of decreased maximum and increased minimum temperatures.

The dynamics responsible for changes within the annual cycle or between different periods have not yet been unequivocally identified. The location of the polar front jet stream and its associations with solar radiation, clouds, snow cover and other atmospheric and surface variables are certainly involved. We believe that the identification of the climate system dynamics responsible for the structure of the annual cycle of daily temperature is a necessary first step in the study of potential human-induced climate change.

1. Introduction

In recent years a large body of research has been directed toward the detection of potential changes that may be taking place within the Earth's climate system. More sensitive

approaches to detection are being made possible by advancements in two key areas: progress in techniques related to modeling the Earth's climate system, and the refinement of quality-controlled climatic data sets that allow intensive investigations of climate change issues. Many detection studies have been based on fingerprint methods (Madden and Ramanathan, 1980; Barnett, 1986). In this procedure, baseline model simulations are completed to provide a point of comparison for later model runs that include increased levels of CO₂ or other greenhouse gases. This allows the models to provide a suite of potential transformations from the present climate state to a new one that is in equilibrium with the changed atmospheric greenhouse-gas content. These templates, or fingerprints, can be used to guide researchers in searching for specific climate change signals within observed records that match the patterns found by the models. These and other hybrid methods of climate change detection (Barnett and Schlesinger, 1987; Wigley and Barnett, 1990) have been unable as yet to detect an unambiguous climate change signal.

One of the greatest weaknesses of detection methodologies in use today is their reliance on monthly or annual averages in conducting fingerprint comparisons. Although these low-frequency statistics are usually the simplest to employ, there is a considerable loss of information when daily data are averaged over longer periods. Specifically, information concerning changes in the shape of the annual cycle is obscured in these procedures.

The use of daily data in climate change studies has been limited in the past by the lack of quality controlled, homogeneous datasets that have a long period of record and dense station distributions over a large spatial domain. It is well known that temperature–time series can be affected by a multitude of operational changes, including time of observation biases (Karl et al., 1986), station moves, instrument replacements, warming caused by urbanization (Karl et al., 1988; Karl and Jones, 1989), or changes in the station's microclimatic environment. Thus, any daily data used in studies of climate change must undergo rigorous quality control procedures before being used.

The purpose of this paper is to demonstrate the utility of long time series of daily data in the understanding of climate change. We are interested in the low-frequency signal of climate change that is contained within the structure of long continuous time series (up to 100 years) of daily data. Data from the Historical Daily Climate Dataset (Robinson, 1993) are used to generate daily annual cycles of maximum and minimum temperatures at six stations in the eastern United States, three in the U.S. Great Plains and one in the southwest U.S. The annual cycle climatologies for these stations are then examined for a recent 30-year interval, and observed changes in the annual cycle of temperature during this century are investigated at three stations. Finally, we speculate as to the potential climate system dynamics at work within the annual cycle and between different periods. We believe that the identification of the climate system dynamics responsible for the shape of the annual cycle of temperature is a necessary first step in the study of potential human-induced climate change.

2. Background

Investigations into temperature changes using the instrumental record have been conducted across the spectrum of spatial scales, from global studies to examinations at single

locations. The study of planetary-scale variations of surface air temperatures is exemplified by those who have considered changes in areally averaged hemispheric temperatures (Jones et al., 1986; Hansen and Lebedeff, 1987; Angell, 1988; Spencer and Christy, 1990). Others have examined changes in temperature and its variability at continental and hemispheric scales (Barnett, 1978; Gutzler et al., 1988; Karl et al., 1991a), while regional temperature changes have been documented over diverse areas of the globe (Diaz, 1986; Karl et al., 1991b; Michaels et al., 1988; Walsh and Chapman, 1990; Kalkstein et al., 1990). Even though temperature variations at a single location are difficult to interpret because of human-induced changes that can affect a given observing site, several studies have documented changes in the temperature regime at this level (e.g., Agee, 1982; Skaggs and Baker, 1989).

In these studies, which concentrate on mean temperatures, as well as in those which explore long term changes in diurnal temperature ranges (Karl et al., 1984; Karl et al., 1987; Robinson, 1992), daily data are generally aggregated into monthly, seasonal or annual means. While these studies have increased our knowledge of climate dynamics, the aggregation of daily climate variables may result in the loss of valuable information. For example, using daily temperatures for a 74-station network in the eastern United States, Guttman and Plantico (1987) compared the published daily normals, interpolated from monthly averages for a 30-year period, to the calculated 30-year average daily temperatures. They found that the published normals were statistically different from the annual cycles created by using observed daily data, and that singularities seemed to exist in the average annual cycles of the observed data at several of the stations. In a follow up study, the same authors found no significant periodicities in the calculated daily annual cycles. Most recently, Ruschy et al. (1991) have discussed large step changes in the annual cycle of the daily temperature range at several stations in Minnesota.

Few additional studies have examined long (> 30 year) time series of observed daily data. Brinkman (1983) examined daily climate variability in Wisconsin with respect to seasonal temperature extremes, and Agee (1982) and Skaggs and Baker (1989) studied climate changes at singular locations. We know of no investigations of the annual cycle of daily temperature that have been completed using a spatially diverse station network possessing a long period of record.

3. Data

In recognition of the need for lengthy and accurate climatological observations for use in global change studies, a unique digital set of daily climatic data for approximately 1300 United States stations has been assembled (Robinson, 1993). This set, known as the Historical Daily Climate Dataset (HDCD), contains observations of precipitation, snowfall, snow on the ground and maximum and minimum temperature. Records extend from 1988 back to the turn of the century for about 400 of the stations, and to the 1920's and 1930's for most others. Similar digital data are unavailable for almost all other U.S. stations before 1948. The quality of all data (each variable for each day of record) is examined, and flags representing the results of the query are incorporated into the set. Included in the set are station observations from every conterminous U.S. state, with the exception of Delaware.

Table 1
Location and elevation of study stations

Station	Latitude	Longitude	Elevation (m)
Belle Glade, Fla.	26°39'	80°38'	6
Burlington, Vt.	44°28'	73°09'	101
Caribou, Maine	46°52'	68°01'	190
Cooperstown, N.Y.	42°42'	74°55'	366
Gothenburg, Nebr.	40°56'	100°10'	783
Grand Island, Nebr.	40°58'	98°19'	566
Washington, Iowa	41°17'	91°41'	230
Willcox, Ariz.	32°18'	109°51'	1273
Winnsboro, S.C.	34°23'	81°06'	171
Woodstock, Md.	39°20'	76°52'	140

Most are cooperative stations or smaller first order sites, which are less likely to be influenced by urbanization than larger first order stations.

For the purposes of this introductory study, ten stations from the HDCD will be examined. These include Caribou, Maine; Burlington, Vt.; Cooperstown, N.Y.; Woodstock, Md.; Winnsboro, S.C.; Belle Glade, Fla.; Grand Island, Nebr.; Gothenburg, Nebr.; Washington, Iowa and Willcox, Ariz. (Table 1). Daily temperature data for the periods of record discussed here are 99% or more complete, and of the available data, less than 0.2% are flagged as suspect at any station. For the purposes of this study, it was not necessary to fill in missing data, and all flagged data are excluded from the analyses.

4. Daily annual cycles

4.1. Maximum and minimum temperatures

Daily annual cycles are derived here from maximum and minimum temperatures by calculating the long-term daily mean for each day of the year. The traditional method used to generate daily means is the application of a spline, fitted to 30-year monthly averages and evaluated for each day. This may suffice for many climate applications, but obscures potentially useful climate information. For example, at Cooperstown, N.Y. (Fig. 1a), extended periods of alternating cold and warm intervals during January and February do not follow the spline fitted data. Deviations that last for extended periods are also common at Gothenburg, Nebr. (Fig. 1b), especially in the maximum temperature annual cycle.

4.2. Temperature range

We have found that the daily temperature range exhibits rapid variations over the course of the mean annual cycle. Temperature range is defined as the difference between mean maximum and mean minimum temperatures on a given day. Major discontinuities in temperature range are observed at many stations. Step changes may occur over periods as short as several days. For example, the mean daily temperature range at Burlington, Vt. falls from




Fig. 1. Annual cycle of daily maximum and minimum temperatures at: (a) Cooperstown, N.Y. and (b) Gothenburg, Nebr. Based on a 30 year mean of daily values (1959–1988) (thin line) and a spline fit applied to mean monthly temperatures during this period (thick line).

approximately 13°C to 8°C over a three day period in early November (Fig. 2a). This is the result of an accelerated decrease in maxima, accompanied by a slight rise in minima for a short period prior to a continuation of their seasonal decline. There is also a step change from 10°C to 13°C over approximately two weeks in the middle of April. This results from an accelerated increase in minima, accompanied by a slight decrease in maxima for a short period prior to a continuation of their seasonal rise. These step changes are quite large, despite the fact that the daily data are averaged over a 30-year period. These discontinuities are also found with the same timing and magnitude at other stations in Vermont (Fig. 2b and c).

An inspection of annual cycles of daily temperature range at stations along the east coast of the United States finds abrupt transitions of varying magnitude and length (Fig. 3). The daily range cycle at Caribou, Maine (Fig. 3a) has the same general characteristics as Burlington (Fig. 2a), including a summer maximum and winter minimum. The autumn decrease in range is not as pronounced as in Vermont, occurring over the course of six weeks. Rather than a single step in spring, Caribou's range first decreases for one to two weeks in early April and then rises to summer levels by early May.

Overall, the daily temperature range is similar at Caribou and Burlington, and only about 2°C degrees less pronounced than at Cooperstown, N.Y. (Fig. 3b), the next station examined in a southward progression along the interior East Coast. The shape of the New York station's annual cycle is quite similar to the Vermont station. This is most apparent in fall, when the discontinuity in range is coincident with that of Burlington. The spring increase in range at Cooperstown is more gradual than in Vermont. It is similar to that observed at Woodstock, Md. (Fig. 3c), although at the Maryland station this rise is rather continuous from an early winter range minimum. The range decrease at Woodstock in autumn begins




Fig. 2. Mean daily annual cycles of temperature range (maximum minus minimum) for three stations in Vermont.

at about the same time as the more northern stations, but is more gradual. A fall maximum in temperature range, approaching that of late spring, is observed at Woodstock, but not further north. The magnitude of the daily temperature range is similar at Woodstock and Cooperstown.

While the magnitude of the annual cycle of daily temperature range is only about 1°C higher at Winnsboro, S.C. (Fig. 3d) than at Woodstock, a very different shape to the cycle is evident. A summer range minimum occurs in South Carolina, with range maxima in the transitional seasons. Winter daily ranges are slightly larger than in summer. No pronounced discontinuities are seen at Winnsboro, the decrease in range from spring to summer being quite gradual, with the fall increase occurring throughout October. It is interesting that the




Fig. 3. Mean daily annual cycles of temperature range (maximum minus minimum) for five stations in the eastern United States.

fall peak in South Carolina is coincident with the sharp downward discontinuity at the northern stations. Such a fall peak is not observed further south at Belle Glade, Fla. (Fig. 3e), rather a gradual increase in daily temperature range is observed from October to late January. At this time, the temperature range levels off until a rather pronounced decrease of more than 2°C starting at the end of April and continuing into the middle of May. The range varies little from June through September. While the annual cycle is less pronounced at Belle Glade, the magnitude of the daily temperature range is similar to that found elsewhere in the eastern U.S.

Distinctive annual cycles of daily temperature range are also observed in other areas of the United States. For example, at Grand Island, Nebr., the flow of the cycle during the year




Fig. 3. (continued).

is smoother than in Eastern regions (Fig. 4a). The shape of the cycle is most similar to Woodstock, Md. in the East. This is interesting, given that these stations are at about the same latitude. In the southwestern U.S., there is a pronounced decrease in the daily temperature range during early July (Fig. 4b). A gradual increase at Willcox, Ariz. follows, progressing to an early November maximum, much as is observed at Winnsboro, S.C. At each of the stations discussed above variations in the annual cycles of both maximum and minimum temperature contribute to the abrupt changes and discontinuities.





Fig. 4. Mean daily annual cycles of temperature range for Grand Island, Nebr. and Willcox, Ariz. Note the vertical scale for 4b has been altered to accommodate the data.




Fig. 5. Differences in mean daily annual cycles of minimum temperature in January through March between 1952–1967 and 1973–1988 (difference equals 73–88 minus 52–67).

4.3. Interdecadal variations

Through time, change in the shape of the daily annual cycle of temperature variables may indicate changing climate system dynamics. Within a single month, large variations are observed in the mean daily differences that would not be distinguishable in monthly data. For instance, at Washington, IA, daily minimum temperatures during one 15 year period (1973–1988) were more than 3°C colder during the first half of January than during the 1952–1967 period (Fig. 5). Opposite conditions were observed during the second half of the month. If one were to calculate the monthly difference, the two anomalies would nearly cancel, leading to the erroneous conclusion of no change. A similar situation can be observed in February, also, but not in March. Small changes in the timing of seasonal warming or cooling can create very large differences in temperature between two sub-monthly periods.

Examinations of the full annual cycle of daily temperatures show unique characteristics within certain decades. For example, mean daily maximum temperatures at Grand Island, Nebr. were warmer in the 1930's compared to the entire 1900–1988 period on approximately 90% of the days of the year (Fig. 6a). The warmth was most pronounced during summer. Daily minima showed similar tendencies, resulting in little change in daily temperature ranges over the course of the year (Fig. 6b).

The situation in the 1930's was different at Burlington, Vt., where mean daily maximum temperatures showed only the slightest tendency towards being cooler than over the entire 1900–1988 period (Fig. 6c). Minima showed some increase in the 1930's at Burlington, combining with the maxima to result in a stronger tendency towards a suppressed daily temperature range during this period (Fig. 6d).




Fig. 6. (a) Differences in mean daily annual cycles of maximum temperature at Grand Island, Nebr. between 1900–1988 and 1930–1939 (difference equals 30–39 minus 00–88). (b) Same as Fig. 6a, except for temperature range. (c) Same as Fig. 6a, except for Burlington, Vt. d. Same as 6a, except for temperature range at Burlington, VT.

5. Discussion

Our efforts to date have concentrated on the creation of the daily dataset and identification of the structure of daily temperature range cycles. It is beyond the scope of this paper to unequivocally identify the dynamics responsible for changes within the annual cycle or between different periods. Here, we briefly hypothesize as to the possible causes of the abrupt transitions observed in the daily annual temperature cycles. Depending on the region, these may include polar front positioning, the location of seasonal highs and lows, or monsoonal dynamics. These circulation features are associated with variations in atmospheric and surface moisture, phenology, clouds and snow cover, all of which influence the

radiation balance, and, ultimately, surface air temperatures. The arrival of the polar front into the northeastern U.S., with an associated increase in cloudiness, may explain the abrupt reduction in daily temperature range in fall. This discontinuity precedes the establishment of a persistent snow cover by several weeks, thus snow can not be invoked as an explanation. At Caribou, Maine, the timing of snow melt in spring is more in line with changes in the temperature range. Maxima may be held close to the freezing point until the snow disappears, potentially explaining the temporary pronounced decrease in the temperature range. The spring “greenup” (Schwartz and Karl, 1990; Schwartz, 1992) in the middle latitudes may influence daily temperature ranges. Perhaps phenologic changes at other times of the year also influence range shifts.

Summer range minima in the southeastern U.S. may be due to the onset of persistent maritime-tropical air masses. With them, atmospheric moisture increases, keeping daily minima higher, and afternoon cloudiness is enhanced, suppressing maxima. This situation is less common during the transitional seasons, leading to an increased range. Lower solar radiation and perhaps increased cloudiness associated with mid-latitude disturbances lower the winter range from the transition season maxima. This situation is more pronounced in the Southwest, with monsoonal clouds and precipitation affecting both maximum and minimum temperatures in summer.

A tendency towards greater continentality of the climate over the central United States in the 1930’s, with associated drier atmospheric and surface conditions, explains increased maxima and minima at Grand Island, Nebr. These conditions were apparently not felt in the northeastern U.S., where a suppressed temperature range (a result of decreased maxima and increased minima) may have resulted from increased cloudiness. In the case of inter-decadal comparisons in Washington, Iowa, the abrupt changes from colder to warmer relative conditions may be associated with changes in the location, intensity, and timing of the establishment of a longwave trough over the eastern U.S.

Work has commenced to increase the spatial coverage of the temporal analyses, as well as to diagnose the causes of variations in the daily temperature cycles. As mentioned previously, no other studies have explored daily temperature cycles at broad spatial dimensions. Only one local study has attempted to diagnose abrupt changes in the annual temperature cycle (Ruschy et al., 1991). Changes observed in Minnesota during the fall and spring were found to be related to solar radiation, cloudiness and snow cover. However, with none of these variables does the timing of changes during the year fit the abrupt changes in daily temperature range. The atmospheric circulation may play a key role in the timing of these temperature changes. Thus, temperatures may be influenced by a combination of these and other forcings, or, perhaps, thresholds are exceeded as these variables change, triggering the abrupt changes in temperature range.

6. Summary and conclusions

Distinctive annual cycles of daily maximum and minimum temperature and temperature range are observed across the United States when averaging daily data for multiple decades. In all cases, distinct structures in the range between maximum and minimum temperatures are observed within the annual cycles. Range changes of 5°C may occur within a week,

despite the fact that the daily data are averaged over many decades. Range changes are most pronounced in the fall and spring. Observed differences in mean daily temperature annual cycles between various portions of this century are found for submonthly periods. While the dynamical analysis of causes for these changes is ongoing, atmospheric and surface climate forcings are clearly implicated.

It is apparent that the use of lengthy time series of daily data identifies important information regarding climate diagnostics and climate change detection that is lost when monthly or longer averages are employed. We are continuing to expand the spatial coverage of the temperature analyses, and are beginning rigorous diagnoses of variations in the mean annual daily cycle climatologies as well as interdecadal variations in these cycles. We also plan to compare observed daily annual cycles with those derived from general circulation model output, for the purpose of detecting a greenhouse-gas induced climate change. If observed changes in annual cycles over the Twentieth Century correspond to differences in annual cycles between control and doubled-CO₂ model climates, a large step will have been taken toward confirming the cause of recent climate change.

Acknowledgements

We wish to thank J. Wright and A. Ellis for assisting with data generation. This research was funded by the U.S. Department of Energy's (DOE) National Institute for Global Environmental Change (NIGEC) through the NIGEC Great Plains Regional Center at the University of Nebraska–Lincoln (DOE Cooperative Agreement No. DE-FC03-90ER61010). Financial support does not constitute an endorsement by DOE of the views expressed in this article.

References

- Agee, E.M., 1982. A diagnosis of twentieth century temperature records at West Lafayette, Indiana. *Climate Change*, 4: 399–418.
- Angell, J.K., 1988. Variations and trends in tropospheric and stratospheric global temperatures, 1958–87. *J. Climate*, 1: 1296–1313.
- Barnett, T.P., 1978. Estimating the variability of surface air temperature in the Northern Hemisphere. *Mon. Weather Rev.*, 106: 1353–1367.
- Barnett, T.P., 1986. Detection of changes in the global tropospheric temperature field induced by greenhouse gases. *J. Geophys. Res.*, 91: 6659–6667.
- Barnett, T.P. and Schlesinger, M.E., 1987. Detecting changes in global climate induced by greenhouse gases. *J. Geophys. Res.*, 92: 14,772–14,780.
- Brinkman, W.A.R., 1983. Variability of temperature in Wisconsin. *Mon. Weather Rev.*, 111: 172–180.
- Diaz, H.F., 1986. An analysis of twentieth century climate fluctuations in northern North America. *J. Clim. Appl. Meteorol.*, 25: 1625–1657.
- Guttman, N.B. and Plantico, M.S., 1987. Climatic temperature normals. *J. Clim. Appl. Meteorol.*, 26: 1428–1435.
- Gutzler, D.S., Rosen, R.D. and Salstein, D.A., 1988. Patterns of interannual variability in the Northern Hemisphere wintertime 850 mb temperature field. *J. Climate*, 1: 949–964.
- Hansen, J. and Lebedeff, S., 1987. Global trends of measured surface air temperature. *J. Geophys. Res.*, 92: 13,345–13,372.

- Jones, P.D., Raper, S.C.B., Bradley, R.S., Diaz, H.F., Kelly, P.M. and Wigley, T.M.L., 1986. Northern Hemisphere surface air temperature variations, 1851–1985. *J. Clim. Appl. Meteorol.*, 25: 1213–1230.
- Kalkstein, L.S., Dunne, P.C. and Vose, R.S., 1990. Detection of climatic change in the western North American Arctic using a synoptic climatological approach. *J. Climate*, 3: 1153–1167.
- Karl, T.R., Kukla, G. and Gavin, J., 1984. Decreasing diurnal temperature range in the United States and Canada from 1941–1980. *J. Clim. Appl. Meteorol.*, 23: 1489–1504.
- Karl, T.R., Williams, C.N., Jr. and Young, P.J., 1986. A model to estimate the time of observation bias associated with monthly mean maximum, minimum and mean temperatures for the United States. *J. Clim. Appl. Meteorol.*, 25: 145–160.
- Karl, T.R., Kukla, G. and Gavin, J., 1987. Recent temperature changes during overcast and clear skies in the United States. *J. Clim. Appl. Meteorol.*, 26: 698–711.
- Karl, T.R., Diaz, H.F. and Kukla, G., 1988. Urbanization: its detection and effect in the United States climate record. *J. Climate*, 1: 1099–1123.
- Karl, T.R., Heim, R.R. and Quayle, R.G., 1991a. The greenhouse effect in central North America: If not now, when? *Science*, 251: 1058–1061.
- Karl, T.R., Kukla, G., Razuvayev, V., Changery, M., Quayle, R.G., Heim, R.R., Jr., Easterling, D. and Fu, C.B., 1991b. Global warming: evidence for asymmetric diurnal temperature change. *Geophys. Res. Lett.*, 18: 2253.
- Karl, T.R. and Jones, P.D., 1989. Urban bias in area-averaged surface air temperature trends. *Bull. Am. Meteorol. Soc.*, 70: 265–270.
- Madden, R.A. and Ramanathan, V., 1980. Detecting climate change due to increasing carbon dioxide. *Science*, 209: 763–768.
- Michaels, P.J., Sappington, D.E. and Stooksbury, D.E., 1988. Anthropogenic warming in North Alaska? *J. Climate*, 1: 942–946.
- Robinson, D.A., 1993. Historical daily climatic data for the United States. In: Preprints of the Eighth Conference on Applied Climatology, Anaheim, CA. *Am. Meteorol. Soc.*, pp. 264–269.
- Robinson, P.J., 1992. Trends in the relationship between monthly and daily temperatures across the United States. *Phys. Geogr.*, 13: 191–205.
- Ruschy, D.L., Baker, D.G. and Skaggs, R.H., 1991. Seasonal variation in daily temperature ranges. *J. Climate*, 4: 1211–1216.
- Schwartz, M.D., 1992. Phenology and springtime surface layer change. *Mon. Weather Rev.*, 120: 2570–2578.
- Schwartz, M.D. and Karl, T.R., 1990. Spring phenology: Nature's experiment to detect the effect of "green-up" on surface maximum temperatures. *Mon. Weather Rev.*, 118: 883–890.
- Skaggs, R.H. and Baker, D.G., 1989. Temperature change in eastern Minnesota. *J. Climate*, 2: 629–630.
- Spencer, R.W., and Christy, J.C., 1990: Precise monitoring of global temperature trends from satellites. *Science*, 247: 1558–1562.
- Walsh, J.E. and Chapman, W.L., 1990. Short-term climatic variability of the Arctic. *J. Climate*, 3: 237–250.
- Wigley, T.M.L. and Barnett, T.P., 1990. Detection of the greenhouse effect in the observations. In: J.T. Houghton, G.J. Jenkins and J.J. Ephraums (Editors), *Climate Change, The IPCC Scientific Assessment*. Cambridge University Press, pp. 243–255.